

# Conformal maps in periodic dynamo flows and in stretch-twist suppression on Riemannian manifolds

by

*L.C. Garcia de Andrade*

Departamento de Física Teórica – IF – Universidade do Estado do Rio de Janeiro-UERJ  
Rua São Francisco Xavier, 524  
Cep 20550-003, Maracanã, Rio de Janeiro, RJ, Brasil  
Electronic mail address: garcia@dft.if.uerj.br

## Abstract

Examples of conformal dynamo maps have been presented earlier [Phys Plasmas **14**(2007)] where fast dynamos in twisted magnetic flux tubes in Riemannian manifolds were obtained. This paper shows that conformal maps, under the Floquet condition, leads to coincidence between exponential stretching or Lyapunov exponent, conformal factor of fast dynamos. Unfolding conformal dynamo maps can be obtained in Riemann-flat manifolds since here, Riemann curvature plays the role of folding. Previously, Oseledts [Geophys Astrophys Fluid Dyn **73** (1993)] has shown that the number of twisted and untwisting orbits in a two torus on a compact Riemannian manifold induces a growth of fast dynamo action. In this paper, the stretching of conformal thin magnetic flux tubes is constrained to vanish, in order to obtain the conformal factor for non-stretching non-dynamos. Since thin flux tube can be considered as a twisted or untwisting two-torus map, it is shown that the untwisting, weakly torsion, and non-stretching conformal torus map cannot support a fast dynamo action, a marginal dynamo being obtained. This is an example of an anti-fast dynamo theorem besides the ones given by Vishik and Klapper and Young [Comm Math Phys **173**(1996)] in ideally high conductive flow. From the Riemann curvature tensor it is shown that new conformal non-dynamo, is actually singular as one approaches the magnetic flux tube axis. Thus conformal map suppresses the stretching directions and twist, leading to the absence of fast dynamo action while Riemann-flat unfolding manifolds favors non-fast dynamos. **Key-words:** anti-dynamo theorems, conformal stretching. **PACS numbers:** 2.40.Hw: differential geometries. 91.25.Cw-dynamo theories.

# I Introduction

An anti-dynamo theorem by Zeldovich [1], shows that planar flows cannot support dynamo action. Together with Cowling [2] theorem they compose the two first anti-dynamo theorems in magnetohydrodynamics (MHD). As it is well-known, the stretching is a fundamental ingredient in fast dynamo [3] action and leads to exponential stretching. This important role played by stretching in the building of dynamo action led Vishik [4] to formulate the first anti-fast-dynamo theorem, which states that no fast dynamo action can be obtained in non-stretching flows, and only fast dynamos could be obtained. More recently Klapper and Young [5] have extended the  $\mathcal{C}^\infty$ -differentiable anti-fast-dynamo theorem to  $\mathcal{C}^2$ . In this paper, following the conformal technique transformation path, so much used in Einstein's general theory pseudo-Riemannian spacetime, a conformal dynamo flow is proposed by considering a generalization of the Ricca's flux tube [6] to the conformal Riemann metric to his thin twisted magnetic flux tube. An interesting aspect of this technique applied to dynamo theory, is that contrary to what happens in Einstein's theory of gravity, here the conformally flat metric is not flat, in the sense that the Riemann curvature tensor components do not all vanish. As it is shown in the paper, Riemann curvature components depend upon the Frenet curvature and torsion of the magnetic flux tube axis. Throughout the paper, to simplify computations the tube is considered as helical [7] which is so commonly used in plasma astrophysics. This implies also that folding plays the role of Riemann curvature and folding in this sense is not always destructive in the anti-dynamo case considered here since, all components of the Riemann curvature tensor are positive and the folding is constructive [3]. In this way though there is no cancellation of magnetic fields, presence of non-stretching even in the twisting case, leads to a slow dynamo or non-dynamo at all. Actually in the first part of the paper an anti-fast dynamo like theorem is obtained forcing the stretching term in non-dynamo equation to vanish. This is mainly due to the fact that, cancellation of magnetic fields are possible in regions of folding or curvature. In tubes, the Riemann curvature concentrates strongly the magnetic fields inside the tubes in order that the cancellation cannot take place. In the second part of the paper one considers that actually that conditions and constraints that led to the non-dynamo are actually too restrictive or demanding and that fast conformal dynamos [8] can be obtained in Riemannian

spaces. Actually Arnold, Zeldovich, Ruzmaikin and Sokoloff [9] and Chicone and Latushkin [10] have previously shown that uniform stretching in flows could lead to fast dynamos in compact Riemannian manifolds. Though in the first part of the paper one addresses the conformal slow dynamos in the zero-resistivity limit, in the second part one considers conformal fast dynamos in diffusive media. Actually diffusion processes in Riemannian manifolds have been already considered by S. Molchanov [11], without addressing the problem of dynamo action. Indeed Soward [12], argued earlier that fast dynamo actions would still be possible in regions where no non-stretching flows would be presented, such as in some curved surfaces. Here one argues that, there is no reason to believe that these surfaces could not be Riemannian. In this paper the well-known stretch, twist and fold method developed by Vainshtein and Zeldovich [13] seems to be effective due to the conformal dynamo map. Actually from the mathematical point of view, Osedeleets [14] has shown that by investigating the fast dynamo Lyapunov exponent on a two-torus map, is the untwisting and twisting number. To resume, due to the frozen magnetic field from the zero resistivity hypothesis, the flow stretching necessary leads to the geometrical stretching of the Riemann metric. It seems that similar to Soward's argument, the Riemann curved surfaces of the tube which are non-stretched, necessarily leads to non-dynamos, dynamo solutions being possible off the surfaces. The paper is organized as follows: In section II the conformal map is shown to lead , under Floquet condition, to the fast dynamo where the conformal Riemann metric factor is the exponential stretching with Lyapunov exponent of chaotic flows. In section III an anti-fast dynamo is proved and the Riemann curvature tensor are computed and stretching instability of the conformal class discussed in section II is obtained. In section III the suppression of stretching and twist by conformal action is shown to lead to a marginal, non-fast-dynamo in compact Riemannian manifold. In section IV a conformal dynamo map in twisted magnetic flux tube is obtained. Discussions and future prospects are presented in section V.

## II Conformal fast dynamo maps on periodic flows

Childress and Gilbert [3] have discussed in detail the suppression of stretching and twist on the flux tubes. In this section a very simple proof is done of the fact that the conformal metric factor is given by the exponential stretching leading to the conformal fast dynamo. In the next two sections, suppression of stretch and twist in turn, is showed to be connect to non-fast dynamos, providing us with a new application of Vishiks and Klapper and Young anti-fast dynamo theorems. In this section one shall basically deal with Cartesian coordinates in three-dimensional Euclidean  $\mathbf{E}^3$ . Let us start by considering the conformal transformation or rescaling  $\Omega$ , in the Cartesian coordinates as

$$x'^i = \Omega^{-1} x^i \quad (\text{II.1})$$

where the indices  $(i, j = 1, 2, 3)$ . From this equations

$$\frac{\partial x'^i}{\partial x^j} = \Omega^{-1} \delta^i_j \quad (\text{II.2})$$

Thus from the invariance of line elements of Riemann differential geometry ,and by the law of transformation of the Riemann metric components  $g_{ij}$  one has

$$g'_{ij} = \Omega^2(\mathbf{x}, t) g_{ij} \quad (\text{II.3})$$

By considering the general expression of the magnetic field variation under the general coordinate transformation  $\mathbf{x}' \rightarrow \mathbf{x}$ , one has to use the Jacobian transformation [3]

$$B'^i = \mathbf{J}^i_j B^j \quad (\text{II.4})$$

where the Jacobian matrix  $\mathbf{J}^{ij}$  is

$$\mathbf{J}^i_j := \frac{\partial x'^i}{\partial x^j} \quad (\text{II.5})$$

Application of these formulas into the conformal transformation of the partial derivative in the gradient

$$(\nabla')_i := \partial'_i = \Omega \nabla_i \quad (\text{II.6})$$

comparision with expression (II.1) helps us to determine the conformal factor  $\Omega$  as

$$\Omega^{-1} \delta^i_j = \mathbf{J}^i_j \quad (\text{II.7})$$

which from (II.4) yields

$$B'_i = \Omega B_i \quad (\text{II.8})$$

when one consider the periodic flows under the Flouquet condition

$$B'_i(\mathbf{x}, t + T) = e^{\gamma(\eta)T} B_i(\mathbf{x}, t) \quad (\text{II.9})$$

which from (II.8) yields

$$\Omega = e^{\gamma T} \quad (\text{II.10})$$

where  $\gamma$  is the growth of magnetic field. Therefore one must conclude that under certain conditions on diffusion, the conformal factor can be associated with exponential chaotic stretching. In the next section one adopts the more simple assumption of zero-resistivity plasmas. Since the dynamo action is basically discovered by solving the self-induction equation, most of the times in Cartesian coordinates, now let us investigate this equation in this coordinates as

$$d_t B'^i = B'^j \partial'_j v^i + \eta \partial'^k \partial'_k B^i \quad (\text{II.11})$$

Now by considering the zero-resistivity ideal plasma, where resistivity  $\eta$  vanishes, one has

$$d_t B'^i = B'^j \partial'_j v^i \quad (\text{II.12})$$

By taking into account that flow vector transform conformally as

$$v'_i = \Omega v_i \quad (\text{II.13})$$

one obtains the following constraint on the omega conformal factor as

$$d_t \Omega + \frac{1}{3}(\mathbf{v} \cdot \nabla) \Omega = 0 \quad (\text{II.14})$$

where one has made use of the unprimed self-induced equation to achieved this equation. This equation is very important even when periodic flows are absent, as we shall see by the end of this section , when the unfolding is connected with the vanishing of the Riemann tensor, or like in geometers jargon, a Riemann-flat manifold. In this case one shall drop the Cartesian coordinates used here and consider the Riemannian tubular coordinates to a, in principle curved Riemannian flux tube. Since we are considering in this section all conformal dynamo maps and leave the anti-dynamo theorems and non-dynamos to the last two sections, let us consider first

the case of flux tubes where the conformal factor  $\Omega$  depends only on the radial coordinate- $r$ . Thus by expression

$$(\mathbf{v} \cdot \nabla) \Omega = 0 \quad (\text{II.15})$$

applied to magnetic flux tube coordinates  $(r, \theta_R, s)$ , yields

$$(\mathbf{v} \cdot \nabla) \Omega = v^r \partial_r \Omega(r) = 0 \quad (\text{II.16})$$

which yields  $v^r = 0$  or  $v_r = 0$  since the Riemann metric of the tube ((see formulas in section II)), is diagonal. Thus this choice of conformal factor is consistent with the usual choice of flux tubes, where the flow is confined inside the tube. This is also the approach in plasma torus called tokamaks by plasma physicists. Actually since the folding in flux tubes maybe be represented by the Riemann curvature tensor, constructive folding that leads to fast dynamos, can be obtained by the vanishing of folding or by vanishing of the Riemann curvature tensor. By the formula, for the Riemann curvature tensor for  $\Omega(r)$  below , one finds that  $R_{r\theta r\theta} := 0$  constraint reduces to a differential equation for the conformal factor as

$$r(\Omega^2)'' + (\Omega^2)' - \frac{1}{2\Omega^2}(\Omega^2)'r = 0 \quad (\text{II.17})$$

which has a very simple particular solution compatible with (II.14) which is  $\Omega(r) = r$ , which yields the fast dynamo conformal thin flux tube Riemannian metric as

$$ds_0^2 = r^2[dr^2 + r^2 d\theta^2] + ds^2 \quad (\text{II.18})$$

In this way a fast conformal dynamo map was obtained in periodic and plasma flows inside Riemann-flat thin tubes. Actually from the incompressible flows condition

$$\nabla \cdot \mathbf{v} = 0 \quad (\text{II.19})$$

and the conformal invariance, one is led to conclude that the equation (II.15), is nothing but the statement that the conformal factor does not change along the streamlines. The same is true to the magnetic field lines which solenoidal vector condition

$$\nabla \cdot \mathbf{B} = 0 \quad (\text{II.20})$$

also leads to

$$(\mathbf{B} \cdot \nabla) \Omega = 0 \quad (\text{II.21})$$

which is relation which expresses the fact that the conformal factor does not vary along magnetic field lines. The fact that these two relations are similar, stems from the fact that the plasma flow is ideal or resistivity free, and the frozen condition which tights the magnetic field lines to the streamlines is given. Note also that magnetic expression, the case of magnetic flux tubes yields that  $B^r = 0$  or that the radial component of the magnetic field vanishes due to the confinement condition on astrophysical plasma flux tubes and plasma torus devices.

### III Conformal non-stretching marginal tube dynamos

This section contains the presentation and a simple proof of the following theorem:

**Theorem 1:** Let us consider a Riemannian compact and curved conformal thin magnetic twisted flux tube  $\mathcal{T}$  embedded in the Euclidean space  $\mathbf{E}^3$ . Thus a class of conformal  $\mathcal{C}^1$ -differentiable, slow dynamos, in diffusive-free manifolds can be obtained by imposing the constraint of vanishing of the stretching term of self-induction magnetic equation. Of course, when one considers the resistivity in dynamo flows, the Klapper-Young's  $\mathcal{C}^2$ -differentiable condition is needed.

**Proof:** By considering the conformal metric to the Riemann-flat thin magnetic flux tube metric

$$ds_0^2 = dr^2 + r^2 d\theta_R^2 + ds^2 \quad (\text{III.22})$$

as

$$dl^2 := \Omega^2(r, s) ds_0^2 = \Omega^2[dr^2 + r^2 d\theta_R^2 + ds^2] \quad (\text{III.23})$$

where the coordinates are  $(r, \theta_R, s)$  and  $\theta(s) = \theta_R - \int \tau(s) ds$ ,  $\tau$  being the Frenet torsion. The conformal factor is given by the real function  $\Omega$  which here we must consider at least  $\mathcal{C}^1$  differentiable. Now let us consider the self-induction equations as

$$d_t \mathbf{B} = (\mathbf{B} \cdot \nabla) \mathbf{v} + \eta \nabla^2 \mathbf{B} \quad (\text{III.24})$$

where the resistivity  $\eta = 0$ , since one is assuming the ideal plasma case. By considering that the magnetic field is strictly confined along and inside the tube, and that the resistivity free case leads to the frozen condition, the flow is also confined on the tube which allows us not to stretch the magnetic flux tube without stretching the magnetic field. Therefore, as one shall demonstrate in this section the non-stretching of the magnetic flow along the tube implies necessarily a geometrical non-stretching along the magnetic field axis direction of the the flux tube. The above confinement hypothesis leads to the vanishing of radial components of magnetic field and flow, thus the stretching term in the conformal metric yields

$$(\mathbf{B} \cdot \nabla) \mathbf{v} = \mathbf{e}_r \left( \frac{\partial_s \Omega}{\Omega} \right) A(r, s) + \mathbf{e}_\theta \left[ \frac{B_\theta}{\Omega r} C(r, s) + \frac{B_s}{\Omega} (\tau_0^{-1} \frac{\partial_s \Omega}{\Omega} v_s + \partial_s v_\theta) \right] - \mathbf{t} \left[ \frac{B_\theta}{\Omega} (v_\theta \frac{\partial_s \Omega}{\Omega}) + \frac{B_s}{\Omega} \tau_0^{-1} v_\theta \frac{\partial_s \Omega}{\Omega} \right] \quad (\text{III.25})$$



where

$$A(r, s) := B_\theta \tau_0^{-1} v_\theta - B_s v_s \quad (\text{III.26})$$

and

$$C(r, s) := -\tau_0^{-1} \partial_s v_\theta + r v_s \frac{\partial_s \Omega}{\Omega} \quad (\text{III.27})$$

where by assuming the vanishing of stretching expression (III.27) and considering the other differential equation from the solenoidal vector condition

$$\nabla \cdot \mathbf{B} = [B_\theta \tau_0^{-1} - r B_s] \partial_s \ln \Omega + \Omega^2 \tau_0^{-1} \partial_s B_\theta = 0 \quad (\text{III.28})$$

A almost trivial solution of this system of PDEs is given simply by

$$\frac{\partial_s \Omega}{\Omega} = 0 \quad (\text{III.29})$$

and

$$\partial_s B_\theta = 0 \quad (\text{III.30})$$

This last expression implies that the tube is untwisting.  $\square$

This simple case and actually all solutions of the self-induction condition for this metric possesses a problem which is the fact that the metric that expands along both directions of the tube, namely the tube cross-section and the toroidal direction along the tube magnetic axis, certainly "explodes" and shall never be a dynamo at all, from the very beginning. To avoid these kind of pathologies, one makes use of the non-uniform metric

$$ds_0^2 = \Omega^2 [dr^2 + r^2 d\theta_R^2] + K^2 ds^2 \quad (\text{III.31})$$

where when the  $K$  is given as above represents a flux tube. Thus by computing the same stretching term above with the new sectionally or piecewise metric (III.31) yields

$$\frac{\partial_r(r\Omega)}{\Omega^2} B_\theta v_\theta = -\frac{B_s}{\Omega K} v_s \partial_r K \quad (\text{III.32})$$

$$\frac{B_\theta}{\Omega K} \partial_s \Omega v_s = -\frac{B_s}{K} (\partial_s B_\theta) + \frac{B_s}{r\Omega} \tau_0^{-1} \partial_s K \quad (\text{III.33})$$

$$\frac{B_\theta}{\Omega} \left( \frac{v_\theta}{K} \right) = -\frac{B_s}{K} [\partial_r K - \tau_0^{-1} \frac{v_s}{\Omega} \partial_s K] \quad (\text{III.34})$$

The solenoidal condition for the magnetic fields is

$$(B_\theta - \tau_0 B_s) \partial_s \ln(K\Omega) = \partial_s B_\theta \quad (\text{III.35})$$

From these expressions one may be able to prove the following theorem.

**Theorem 2:** A piecewise conformal flux tube geometry leads to a non-stretching tube, obtained from a non-stretching flow. This is a particular application of Vishik's anti-fast dynamo theorem.

**Proof:** Again untwisting solution  $\partial_s B_\theta = 0$ , now reduces the equations (II.16) through (II.19) leads to

$$\partial_s K = 0 \quad (\text{III.36})$$

which represents the non-stretching along the toroidal direction, and

$$\partial_r K = 0 \quad (\text{III.37})$$

from (II.20). These results shows that the tube is geometrically non-stretching along the toroidal and poloidal directions.  $\square$  A corollary can be obtained from this theorem as:

**Corollary:** The Riemannian metric of the conformal non-dynamo map is given by:

$$ds_0^2 = \left(\frac{\Omega_0}{r}\right)^2 [dr^2 + r^2 d\theta_R^2] + K_0^2 ds^2 \quad (\text{III.38})$$

where  $K_0$  is constant. As we shall see below ,when  $K_0 = 1$  one is left with a conformal non-stretched thin magnetic flux tube.

**Proof:** From the expression (II.16) one obtains

$$\partial_r(\Omega r) = 0 \quad (\text{III.39})$$

which yields  $\Omega = \frac{\Omega_0}{r}$  and the conformal Riemann metric (III.22). In the next section, one shall compute the effects on Riemann curvature folding in the conformal metric.

## IV Curvature-folding and suppression of stretch-twist in conformal marginal dynamos

Since the Riemann curvature is not in principle connected to the stretching of the tube but to its folding, one is able to compute the curvature of the tube as Let us now consider the Riemann metric of a flux tube in curvilinear coordinates  $(r, \theta_R, s)$ , where  $\theta = \theta_R - \int \tau(s) ds$ . Here, accordingly to Ricca's flux tube metric,  $K^2 = (1 - \kappa r \cos \theta)^2$ . This expression contributes to the Riemann curvature components. These can be easily computed with the tensor computer package in terms of Riemann components. Adapting the general relativity (GR) tensor package to three-dimensions, yields the following curvature components expressions

$$R_{1313} = R_{rsrs} = -\frac{1}{4K^2}[2K^2\partial_r D(r, s) - D^2] = -\frac{1}{2}\frac{K^4}{r^2} = -\frac{1}{2}r^2\kappa^4\cos^2\theta \quad (\text{IV.40})$$

$$R_{2323} = R_{\theta s \theta s} = -\frac{r}{2}D(r, s) = -K^2 \quad (\text{IV.41})$$

where  $D := \partial_r K^2$ . For thin tubes,  $K^2(r, s) \approx 1$  ( $r \approx 0$ ), these Riemann curvature components reduce to

$$R_{1313} = R_{rsrs} = -\frac{1}{r^2} \quad (\text{IV.42})$$

Let us now compute the Riemann curvature components of the

$$R_{1313} = R_{rsrs} = -\frac{1}{4K^2}[2K^2\partial_r D(r, s) - D^2] = -\frac{1}{2}\frac{K^4}{r^2} = -\frac{1}{2}r^2\kappa^4\cos^2\theta \quad (\text{IV.43})$$

Now let us compute the Riemann tensor for metric (III.22). Since now most of the Riemann components involve the derivatives with respect to toroidal coordinate-s and derivatives of constant  $K'_0$ , the only non-vanishing component of the Riemann metric is

$$R_{1212} = R_{r\theta r\theta} = -\frac{r}{4}[r(\Omega^2)'' + (\Omega^2)' - \frac{1}{2\Omega^2}(\Omega^2)'r] \quad (\text{IV.44})$$

which from (III.22) yields

$$R_{1212} = -\frac{3\Omega_0^2}{r^2} \quad (\text{IV.45})$$

Thus one notices that the Riemann curvature is singular along the non-dynamo flux tube magnetic axis. Now let us prove the following theorem, which is a stronger version of the previous anti-dynamo theorem for flux tubes:

**Theorem 3:** The non-stretching, untwisted conformal dynamo two-torus map on a compact Riemannian manifold, is unstable and leads to a marginal non-fast dynamo, along the Riemann metric of conformally thin untwisted magnetic flux tube. This can be considered as a suppression of the stretching directions by conformal action.

**Proof:** Let us now analyse the LHS of the self-induction magnetic equation

$$d_t \mathbf{B} = d_t(h^2 B_\theta \mathbf{e}_\theta + h^3 B_s \mathbf{t}) = d_t[(r^{-1} \Omega^{-2} B_\theta \dot{\Omega} + r^{-1} \Omega^{-1} \dot{B}_\theta) \mathbf{e}_\theta + r^{-1} \Omega^{-1} \dot{B}_\theta \dot{\mathbf{e}}_\theta - B_s \dot{\mathbf{t}} + \dot{B}_s \mathbf{t}] \quad (\text{IV.46})$$

making use of the following definitions of the poloidal and toroidal components of the magnetic fields as

$$B_\theta = B_\theta^0 e^{\gamma t} \quad (\text{IV.47})$$

and

$$B_s = B_s^0 e^{\gamma t} \quad (\text{IV.48})$$

where superscript-0 index refers to stationary quantities. By making use of the identities on the relations between the two frames , of tube and Frenet frame

$$\dot{\mathbf{e}}_\theta = -\omega_0 \mathbf{e}_r - \sin\theta \tau_0^2 \mathbf{t} \quad (\text{IV.49})$$

in the helical tube. This computation yields three scalar equations along the tube frame  $(\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{t})$  transformed to the Frenet frame with the help of equations

$$\dot{\mathbf{t}} = [\kappa' \mathbf{b} - \kappa \tau \mathbf{n}] \quad (\text{IV.50})$$

$$\dot{\mathbf{n}} = \kappa \tau \mathbf{t} \quad (\text{IV.51})$$

$$\dot{\mathbf{b}} = -\kappa' \mathbf{t} \quad (\text{IV.52})$$

Together with the flow derivative

$$\dot{\mathbf{t}} = \partial_t \mathbf{t} + (\vec{v} \cdot \nabla) \mathbf{t} \quad (\text{IV.53})$$

one obtains

$$(\Omega r)^{-1} [(\frac{\dot{\Omega}}{\Omega} - \gamma) \sin\theta - \omega_0 \cos\theta] B_\theta + \tau_0^2 B_s = 0 \quad (\text{IV.54})$$

$$(\Omega r)^{-1} [(\frac{\dot{\Omega}}{\Omega} - \gamma) \cos\theta - \omega_0 \sin\theta] B_\theta = 0 \quad (\text{IV.55})$$

$$(\Omega r)^{-1} B_\theta \tau_0^2 \sin\theta - \gamma B_s = 0 \quad (\text{IV.56})$$

By applying the hypothesis of weak torsion  $\tau_0$ , which is a very reasonable hypothesis for astrophysical plasma and solar tubes, reduces the three scalar dynamo equations to

$$(\Omega r)^{-1}[(\frac{\dot{\Omega}}{\Omega} - \gamma)\sin\theta - \omega_0\cos\theta]B_\theta = 0 \quad (\text{IV.57})$$

$$\gamma B_s = 0 \quad (\text{IV.58})$$

Though the equations (IV.43) and (IV.40) together in the absence of twist  $\omega_0 = 0$  yields

$$\Omega = \frac{\Omega_0}{r} e^{\gamma t} \quad (\text{IV.59})$$

and  $\gamma = 0$ . Therefore, though equation (IV.43) yields an apparent Lyapunov exponent and consequently a dynamo action, the other equation (IV.44) shows that only a marginal dynamo is found where the magnetic field does not grow in time, and a fast dynamo is not possible, which is the purpose of the demonstration of anti-fast-dynamo theorems.  $\square$

Note that the constraint of untwisting of the flux tube is not really necessary to obtain the non-fast dynamo action of the conformally transformed thin tube, and was only introduced to show that the Lyapunov exponent of fast dynamo action appears into the discussion.

## V Conclusions

Vishik's anti-fast dynamo theorem further generalised by Klapper and Yound is applied here to conformal Riemann metric of thin flux tubes to show that actually even when the tube is allowed to evolve in time no fast dynamo action is obtained under the non-stretching constraint. To recognize when a map or magnetic flow does represent a dynamo or non-dynamo represents an important issue in plasma physics and astrophysics and deserves further investigation. Earlier the role of the curvature, either Gaussian or Riemannian, on the existence of fast dynamos was considered by Chicone-Latushkin [10] expression for  $\gamma$  in the fast dynamo obtained from geodesic flows in Anosov spaces of negative constant Riemannian curvature

$$\gamma = \frac{1}{2}[-\eta(1 + \kappa^2) + \sqrt{\eta^2(1 - \kappa^2)^2 - 4\kappa}] \quad (\text{V.60})$$

where  $\kappa$  is the Gaussian curvature. This expression shows that when this Gaussian curvature of the two-dimensional manifold vanishes, the expression for  $\gamma$  vanishes as well. This shows that in the absence of curvature a marginal non-fast dynamo is obtained. However seems that Chicone-Latushkin result does not apply to the conformal Riemannian non-stretched tubes discussed here. Curvature in certain sense seems to be more connected to folding dynamo processing than to stretching itself. Actually in cases investigated in this paper, stretching is deeply connected to curvature through term  $K(r, s)$  in Ricca's metric. This leads us to conclude that the curvature is actually connected to stretching and folding. Since in the case considered in this section, twist, curvature and stretching are almost neglected the result seems to be physically correct. The theorems discussed here may be extended to include diffusive plasma flows, for example. Note that since, according to Parker [15], the tube twisting increases by radial or cross-section expansion, the above conformal factor  $\frac{\Omega_0^2}{r^2}$ , is compatible with the untwisting hypothesis used in the proofs of above anti-fast-dynamo theorems. This happens because when the radial coordinate increases, the conformal factor acts as "damping" for this action making the tube decrease actually this cross-section as a compression and giving rise to tube untwisting. Another important observation done by Friedlander and Vishik [16], tells us that the simple geometries of magnetic field lines in steady flows it seems unlikely that fast dynamo action could take place. A similar statement has earlier been done by Arnold et al [9], which showed that magnetic Reynolds number  $R_m^{-1} = 0$  steady flows, only give rise

to non-exponential growth of magnetic field and therefore to no fast dynamo action. One notes that it is more difficult to handle the dynamo problem in non-Euclidean spaces, since in this case, the frames evolve dynamically in time as well. Of course introduction of turbulence and mean field magnetohydrodynamics [17], introduce complications in both cases. Vishik's has also shown that the fast dynamo action can be proved for smooth flows in stretching and unstretching directions. Following this idea, in this paper one has shown that it is possible to suppress the stretching directions by the action of the conformal map. In our example, the flux tube toroidal direction is non-stretching while, the poloidal stretching directions are suppressed by conformal map, yielding a anti-fast-dynamo theorem. Fast conformal dynamos in periodic flows in diffusion plasmas, and in the case of zero resistivity plasmas, showing that unfolding Riemann-flat manifolds favors the dynamo plasmas in flux tube while the non-stretching, untwisting tubes favors non-dynamos or marginal ones, or yet in the sense of anti-fast dynamo theorems a non-fast dynamo. Torus fast plasma dynamos seems to be less favoured, than the plasma dynamo cylinders [18]. Jean Luc Thiffeault [19] has recently called my attention to a paper where The Riemann curvature tensor also vanishes as a constraint to two dimensional diffusion in Riemannian manifolds. More realistic fast dynamo picture than the one given here, was recently given by Gilbert [20] which considers chaotic stretch-shearing flows with strong folding and cat maps in torus as well. Strong folding Riemann curvature dynamos could be investigated elsewhere.

## VI Acknowledgements

I am deeply grateful to Andrew Soward, Dmitry Sokoloff, Jean Luc Thiffeault, Andrew D. Gilbert and Renzo Ricca for their extremely kind attention and discussions on the subject of this paper. I would like to dedicate this paper to the 65<sup>th</sup> anniversary of Professor Andrew Soward, which did so much for our understanding of dynamo theory. Thanks are also due to I thank financial supports from Universidade do Estado do Rio de Janeiro (UERJ) and CNPq (Brazilian Ministry of Science and Technology).

## References

- [1] Ya B Zeldovich, JETP Phys **4**, 460 (1957)
- [2] S. Cowling , Magnetohydrodynamics (1964) Oxford.
- [3] S. Childress, A. Gilbert, Stretch, Twist and Fold: The Fast Dynamo (1996),Springer, Berlin.
- [4] M. Vishik, Sov. Phys. Dokl. **33**, 192 (1988).
- [5] I Klapper and L Young , Comm Math Phys. **173** 623 (1995).
- [6] R. Ricca, Solar Physics **172** (1997),241.
- [7] I Klapper and D Longcope, Evolution Equations of Thin Twisted Flux Tubes, in "Workshop on Stellar Dynamos ASP conference series, (1999) M. Nunez and Ferriz-Mas eds.
- [8] L. C. Garcia de Andrade, Physics of Plasmas 14 (2007). L. C. Garcia de Andrade, Non-holonomic dynamo filaments as Arnolds map in Riemannian space, Astronomical notes (2008) in press.
- [9] V. Arnold, Ya B. Zeldovich, A. Ruzmaikin and D.D. Sokoloff, JETP 81 ,n. 6, 2052 (1981). V. Arnold, Appl Math and Mech **36**, 236 (1972).
- [10] C. Chicone, Yu Latushkin and Montgomery, Comm. in Mathematical Physics (1995). C. Chicone and Yu Latushkin, Proc of the American Mathematical Society 125,N. 11,3391 (1997).
- [11] S Molchanov, Russian mathematical surveys (1978).
- [12] A M Soward, Geophys Astrophys Fluid Dyn **53**,81 (1990).
- [13] S.I. Vainshtein, Ya B Zeldovich, Sov Phys Usp 15 ,159 (1972).
- [14] V. Oseledets, Geophysical and Astrophysics Fluid Dynamics **73** 331 (1993).
- [15] E. N. Parker, Cosmical Magnetism, (1980) Oxford University Press.



- [16] S. Friedlander, M. Vishik, Chaos 1(2),198 (1991).
- [17] F. Krause and K-H. Rädler, Mean field Magnetohydrodynamics, (1980) Pergamon Press.
- [18] Z Wang, V Pariev, C Barnes and D Barnes, Phys Plasmas **9**,5 (2002).
- [19] J L Thiffeault and A H Boozer, Chaos **11**,28 (2001).
- [20] A D Gilbert, Proc. Roy Soc. London A, Mathematical Phys **443**, 585 (1993).